

Patient Name: _____

Patient DOB: _____

DSM-5 Criteria for Opioid Use Disorder

OUD is demonstrated by at least 2 out of the 11 criteria below occurring within a year. Severity of OUD is determined based on the number of criteria met

Check any that apply	
<input type="checkbox"/>	Taking opioids in larger amounts or over a longer period of time than intended
<input type="checkbox"/>	Having a persistent desire or unsuccessful attempts to reduce or control opioid use
<input type="checkbox"/>	Spending excess time obtaining, using, or recovering from opioids
<input type="checkbox"/>	Craving opioids
<input type="checkbox"/>	Continued opioid use causing inability to fulfill work, home, or school responsibilities
<input type="checkbox"/>	Continuing opioid use despite having persistent social or interpersonal problems
<input type="checkbox"/>	Lack of involvement in social, occupation, or recreation activities
<input type="checkbox"/>	Using opioids in physically hazardous situations
<input type="checkbox"/>	Continuing opioid use in spite of awareness of persistent physical or psychological problems a) A need for markedly increased amounts of opioids to achieve intoxication or desired effect b) Markedly diminished effect with continued use of the same amount of an opioid
<input type="checkbox"/>	Exhibiting withdrawal symptoms, as manifested by either of the following: a) The characteristic opioid withdrawal syndrome b) The same (or a closely related) substance are taken to relieve or avoid withdrawal symptoms

Total Number of Boxes Checked: _____

Mild: 2-3 criteria **Moderate:** 4-5 criteria **Severe:** greater than or equal to 6 criteria